

LITERATURE CITED

- Alber, M. 2002. A conceptual model of estuarine inflow management. *Estuaries*, 25 (6B): 1246–61.
- Alden, P., R. Cech and G. Nelson. 1998. National Audubon Society Field Guide to Florida. Knopf, New York, NY.
- Allen, R.P. 1942. The Roseate Spoonbill: research report 2. National Audubon Society, New York, 142 ff.
- Augustin, N.H., D.L. Borchers, E.D. Clarke, S.T. Buckland and M. Walsh. 1998. Spatiotemporal modeling for the annual egg production method of stock assessment using generalized additive models. *Canadian Journal of Fisheries and Aquatic Sciences*, 55:2608–21.
- Bancroft, G.T., A.M. Strong, R.J. Sawicki, W. Hoffman and S.D. Jewell. 1994. Relationships among Wading Bird Forage Patterns, Colony Locations, and Hydrology in the Everglades. In *Everglades: The Ecosystem and Its Restoration*, ed. S.M. Davis and J.C. Ogden. St. Lucie Press, Delray Beach, FL. 615–58 pp.
- Barnes, Ferland and Associates (BFA), Inc. 2004. Seagrasses, Associated Fauna and Faunal Habitat Requirements Documentation and Analysis: Final Task 7 Report. Prepared for the South Florida Water Management District, West Palm Beach, FL.
- Battelle. 2004. Literature Review of Salinity Effects on Seagrasses in Florida Bay. Prepared for the South Florida Water Management District, West Palm Beach, FL.
- Beccue, S. 1999. Endangered Species of Everglades National Park. Everglades National Park, Homestead, FL. Available from: <http://www.nps.gov/ever/eco/danger.htm>
- Beeler, I.E. and T.J. O'Shea. 1988. Distribution and Mortality of the West Indian Manatee (*Trichechus manatus*) in the Southeastern United States: A Compilation and Review of Recent Information: NTIS #PB88-207980/AS (two volumes). National Technical Information Service, Springfield, VA, 613 ff.
- Bjork, R. 1996. Roseate Spoonbill. In *Rare and Endangered Biota of Florida* (volume 5): Birds. University Press of Florida, Gainesville, FL.
- Boesch, D.F., N.E. Armstrong, C.F. D'Elia, N.G. Maynard, H.W. Paerl and S.W. Williams. 1993 (September). Deterioration of the Florida Bay Ecosystem: An Evaluation of the Scientific Evidence. Report to the Interagency Working Group on Florida Bay. Everglades National Park, Homestead, FL.

- Boström, C. and E. Bonsdorff. 2000. Zoobenthic community establishment and habitat complexity: the importance of seagrass shoot density, morphology and physical disturbance for faunal recruitment. *Marine Ecology Progress Series*, 205:123–38.
- Boyer, J. and B. Keller. 2003 (April). Nutrient Dynamics. In A Synthesis of Research on Florida Bay, ed. W. Nuttle, J. Hunt and M. Robblee. Compiled for the Florida Bay Science Program, Science Oversight Panel, U.S. Geological Survey, Florida Caribbean Science Center, Gainesville, FL.
- Boyer, J.N., J.W. Fourqurean and R.D. Jones. 1999. Seasonal and long-term trends in water quality of Florida Bay (1987–97). *Estuaries*, 22:417–30.
- Boyer, J.N., P. Sterling and R.D. Jones. 2000. Maximizing information from estuarine and coastal water quality monitoring networks by diverse visualization approaches. *Estuarine, Coastal and Shelf Science*, 50: 39–48.
- Boyer, J.N. and R. D. Jones. 2003. An Integrated Surface Water Quality Monitoring Program for the South Florida Coastal Waters. FY2002 Cumulative Report Prepared for the South Florida Water Management District, West Palm Beach, FL (contract C-10244-A2). Southeast Environmental Research Center, Florida International University, Miami, FL, 67 pp. Available from: <http://serc.fiu.edu/wqmnetwork/>
- Boyer, J.N. and R.D. Jones. 2001. Long-Term Trends in the Water Quality of Florida Bay (1989–2000). Proceedings from the Florida Bay Science Conference, Key Largo, FL (April 23–26, 2001), 65–66 pp.
- Boyer, J.N., J.W. Fourqurean and R.D. Jones. 1997. Spatial characterization of water quality in Florida Bay and Whitewater Bay by multivariate analysis: zones of similar influence (ZSI). *Estuaries*, 20:743–58.
- Brand, L. 2002. The Transport of Terrestrial Nutrients to South Florida Coastal Waters. In The Everglades, Florida Bay, and Coral Reefs of the Florida Keys, ed. J.W. Porter and K.G. Porter. CRC Press, Boca Raton, FL, 361–413 pp.
- Brewster-Wingard, G.L. and S.E. Ishman. 1999. Historical trends in salinity and substrate in central Florida Bay: a paleoecological reconstruction using modern analogue data. *Estuaries*, 22(2B):369–83.
- Brewster-Wingard, G.L., J.R. Stone and C.W. Holmes. 2001. Molluscan faunal distribution in Florida Bay, past and present: an integration of down-core and modern data. *Bulletins of American Paleontology* (Paleoecological Studies of South Florida, ed. B.R. Wardlaw) 361:199–232.
- Browder, J.A. and D. Moore. 1981. A New Approach to Determining the Quantitative Relationship between Fishery Production and the Flow of Fresh Water to Estuaries. In Proceedings of the National Symposium on Freshwater Inflow to Estuaries, Volume 1, ed. R.D. Cross and D.L. Williams. U.S. Fish and Wildlife Service: U.S. Department of the Interior, Washington, D.C.

- Browder, J.A., Z. Zein-Eldin, M.M. Criales, M.B. Robblee, S. Wong, T.L. Jackson and D. Johnson. 2002. Dynamics of pink shrimp (*Farfantepenaeus duorarum*) recruitment potential in relation to salinity and temperature in Florida Bay. *Estuaries*, 25:1355–71.
- Browder, J.A., P.J. Gleason and D.R. Swift. 1994. Periphyton in the Everglades: Spatial Variation, Environmental Correlates, and Ecological Implications. In *Everglades: The Ecosystem and Its Restoration*, ed. S.M. Davis and J.C. Ogden. St. Lucie Press, Delray Beach, FL.
- Browder, J.A., V.R. Restrepo, J.K. Rice, M.B. Robblee and Z. Zein-Eldin. 1999. Environmental influences on potential recruitment of pink shrimp (*penaeus duorarum*) from Florida Bay nursery grounds. *Estuaries*, 22:484–99.
- Butler, M.J. IV, J.H. Hunt, W.F. Herrnkind, M.J. Childress, R. Bertlesen, W. Sharp, T. Matthews, J.M. Field and H.G. Marshall. 1995. Cascading disturbances in Florida Bay, USA: cyanobacterial bloom, sponge mortality, and implications for juvenile spiny lobsters *Panulirus argus*. *Marine Ecology Progress Series*, 129:119–25.
- Carlson, P.R., M.J. Durako, T.R. Barber, L.A. Yarbro, Y. DeLama and B. Hedin. 1990a. Catastrophic Mortality of the Seagrass *Thalassia testudinum*. In the FDER Florida Bay Annual Synthesis: Seagrass Ecology Report. Florida Department of Environmental Regulation, Office of Coastal Zone Management, Tallahassee, FL, 54 pp.
- Carlson, P.R., M.J. Durako, T.R. Barber, L.A. Yarbro, Y. DeLama and B. Hedin. 1990b. Investigations on the Causes, Extent, and Characteristics of a Seagrass Die-Back in Florida Bay. In the FDNR Annual Performance Report. Florida Department of Natural Resources, St. Petersburg, FL.
- Castillo-Rivera, M., M. Montiel. L. Sanvicente Añorve, and R. Zárate. 2005. Spatial, seasonal and diel distribution patterns of two species of mojarra (Pisces: Gerreidae) in a Mexican tropical coastal lagoon. *Journal of Applied Ichthyology* 21: 498-503.
- Castillo-Rivera, M., J.A. Zavala-Hurtado and R. Zárate. 2002. Exploration of spatial and temporal patterns of fish diversity and composition in a tropical estuarine system of Mexico. *Reviews in Fish Biology and Fisheries*, 12:167–77.
- Childers, D.L., J.N. Boyer, S.E. Davis III, C.J. Madden, D.T. Rudnick and F.H. Sklar. 2005. Nutrient concentration patterns in the oligotrophic “upside-down” estuaries of the Florida Everglades. *Limnology & Oceanography*, in press.
- Cianelli, L., K-S. Chan, K.M. Bailey and N.C. Stenseth. 2004. Nonadditive effects of the environment on the survival of a large marine fish population. *Ecology*, 85:3418–27.
- Clarke E.D., L.B. Spear, M.L. McCracken, F.F.C. Marques, D.L. Borchers, S.T. Buckland and D.G. Ainley. 2003. Validating the use of generalized additive models

- and at-sea surveys to estimate size and temporal trends of seabird populations. *Journal of Applied Ecology*, 40:278–92.
- Cohen, A.D. and T.D. Davies. 1989. Petrographic/botanical composition and significance of the peat deposits of Florida Bay. *Bulletin of Marine Science*, 44(1): 515–16 (abstract).
- Cohen, A.D. and W. Spackman. 1972. Methods in peat petrology and their application to reconstruction of paleoenvironments. *Geological Society of America Bulletin*, 83(1):129–42.
- Committee on Restoration of the Greater Everglades Ecosystem. 2002. Florida Bay Research Programs and their Relation to the CERP. National Academy Press, Washington, D.C., 54 pp. <http://books.nap.edu/books/0309084911/html/index.html>
- Corbett, D.R., J. Chatton, W. Burnett, K. Dillon, C. Rutkowski and J.W. Fourqurean. 1999. Patterns of groundwater discharge into Florida Bay. *Limnology and Oceanography*, 44(4): 1045–55
- Corliss, C.J. 1953. Building the Overseas Railroad to Key West. *Tequesta*, 13:3–21 pp.
- Cosby, B.J., W.K. Nuttle and J.N. Fourqurean. 1999. FATHOM: Flux Accounting and Tidal Hydrology at the Ocean Margin: Model Description and Initial Application to Florida Bay. Report to the Florida Bay Project Management Committee (PMC) and the Everglades National Park (ENP), National Park Service. University of Virginia, Department of Environmental Sciences, Charlottesville, VA.
- Cox, J.R., M. Krautz, M. MacLaughlin and T. Gilbert. 1994. Closing the Gaps in Florida's Wildlife Habitat Conservation System. Florida Fish and Wildlife Conservation Commission, Tallahassee, FL.
- Craighead, F.C. 1971. The Trees of South Florida (volume 1): The Natural Environments and Their Succession. University of Miami Press, Coral Gables, FL.
- Cronin, T.M., C.W. Holmes, G.L. Brewster-Wingard, S.E. Ishman, H.J. Dowsett, D. Keyser and N. Waibel. 2001. Historical trends in epiphytal ostracodes from Florida Bay: implication for seagrass and macrobenthic algal variability. In *Paleoecological Studies of South Florida: Bulletins of American Paleontology* (ed. B.R. Wardlaw), 361:159–98.
- Davis, G.E. and J.W. Dodrill. 1989. Recreational fishery and population dynamics of spiny lobsters, *Panulirus argus*, in Florida Bay, Everglades National Park 1977–1980. *Bulletin of Marine Science*, 44:78–88.
- Davis, J.H. 1940. The Ecology and Geologic Role of Mangroves in Florida. Publication 517, Papers from Tortugas Laboratory, Carnegie Institute, Washington, D.C., 32:303–412.

- Davis, J.H. 1943. The natural features of southern Florida, especially the vegetation and the Everglades. *Florida Geological Survey Bulletin*, 25:311.
- Davis, S. 2004 (January). Everglades Mangrove Estuaries Conceptual Model, Appendix A.3. In Comprehensive Everglades Restoration Plan, Part 1: Monitoring and Supporting Research. United States Army Corps of Engineers, Jacksonville, FL, and South Florida Water Management District, West Palm Beach, FL, A53–78 pp.
- Davis, S.M. and J.C. Ogden. 1994. Toward Ecosystem Restoration. In *Everglades: The Ecosystem and Its Restoration*, ed. S.M. Davis and J.C. Ogden. St. Lucie Press, Delray Beach, FL, 769–96 pp.
- Davis, S.M., D.L. Childers, J.J. Lorenz and T.E. Hopkins. 2005. A conceptual model of ecological interactions in the mangrove estuaries of the Florida Everglades. *Wetlands*, in press.
- Dawes, C.J., D. Hanisak and W.J. Kenworthy. 1995. Seagrass biodiversity in the Indian River Lagoon. *Bulletin of Marine Science*, 57(1):59–66.
- Day Jr, J.W., C.A.S. Hall, W.M. Kemp and A. Yanez-Arancibia. 1989. *Estuarine Ecology*. John Wiley and Sons, Inc. 576 pp.
- DeMaria, Karen. 1996. Changes in the Florida Keys Marine Ecosystem Based upon Interviews with Experienced Residents. The Nature Conservancy, Washington, D.C.
- Deuver, M.J., J.F. Meeder, L.C. Meeder and J.M. McCollom. 1994. The Climate of South Florida and Its Role in Shaping the Everglades Ecosystem. In *Everglades: The Ecosystem and Its Restoration*, ed. S.M. Davis and J.C. Ogden. St. Lucie Press, Delray Beach, FL, 225–48 pp.
- Doering, P.H., R.H. Chamberlain and D.E. Haunert. 2002. Using submerged aquatic vegetation to establish minimum and maximum freshwater inflows to the Caloosahatchee Estuary, Florida. *Estuaries* 25:1343–54.
- Duffy, K.C. and D.M. Baltz. 1998. Comparison of fish assemblages associated with native and exotic submerged macrophytes in the Lake Ponchartrain estuary, USA. *Journal of Experimental Marine Biology and Ecology* 223:199–221.
- Dunson, W.A. 1982. Salinity relations of crocodiles in Florida Bay, USA. *Copeia* 1982(2):374–85.
- Dunton, K. H. 1996. Photosynthetic production and biomass of the subtropical seagrass *Halodule wrightii* along an estuarine gradient. *Estuaries* 19:436–47.
- Dunton, K.H. 1990. Production ecology of *Ruppia maritima* and *Halodule wrightii* in two subtropical estuaries. *Journal of Experimental Marine Biology and Ecology* 143:147–64.

- Durako, M.D. and K.M. Kuss. 1994. Effects of *Labyrinthula* infection on the photosynthetic capacity of *Thalassia testudinum*. *Bulletin of Marine Science*, 54:727–32.
- Durako, M.J., J.C. Zieman and M. Robblee. 2003. Seagrass Ecology. In A Synthesis of Research on Florida Bay, ed. W. Nuttle, J. Hunt and M. Robblee. Compiled for the Florida Bay Science Program, Science Oversight Panel, U.S. Geological Survey, Florida Caribbean Science Center, Gainesville, FL.
- Durako, M.J., M.O. Hall and M. Merello. 2002. Patterns of Change in the Seagrass-Dominated Florida Bay Hydroscape. In The Everglades, Florida Bay, and Coral Reefs of the Florida Keys: An Ecosystem Sourcebook, ed. J.W. Porter and K.G. Porter. CRC Press, Boca Raton, FL, 523–37 pp.
- Dwyer, G.D. and T.M. Cronin. 2001. Ostracode shell chemistry as a paleosalinity proxy in Florida Bay. *Bulletin of American Paleontology*, 361: 249–76.
- Egler, F.E. 1952. Southeast saline Everglades vegetation and its management. *Geobotanica*, 3(4–5):213–65.
- Ehrhardt, N.M. and C.M. Legault. 1999. Pink shrimp, *Farfantepenaeus duorarum*, recruitment variability as an indicator of Florida Bay dynamics. *Estuaries*, 22:471–83.
- Enfield, D.B., A.M. Mestas-Núñez and P. Trimble. 2001. The Atlantic multidecadal oscillation and its relation to rainfall and river flows in the continental U.S. *Geophysical Research Letters*, 28(10):2077–80.
- Enos, P. and R.D Perkins. 1979. Evolution of Florida Bay from island stratigraphy. *Geological Society of America Bulletin*, 90(1):59–83.
- Environmental Consulting and Technology (ECT), Inc. 2005. Project Team: Marshall, F.M., B. Cosby and W. Nuttle. FATHOM Enhancements and Implementation to Support Development of MFL for Florida Bay: Final Report (contract C-C-15975-WO05-05). September 2005. Prepared for the South Florida Water Management District, West Palm Beach, FL, by Environmental Consulting and Technology, Inc.. New Smyrna Beach, FL. 80 pp.
- Estévez, E.D. 2000. A Review and Application of Literature Concerning Freshwater Flow Management in Riverine Estuaries: Technical Report 680. Prepared for the South Florida Water Management District, West Palm Beach, FL. Center for Coastal and Tropical Ecology, Mote Marine Laboratory, Sarasota, FL.
- Estuarine Research Federation. 2002. Dedicated Issue of *Estuaries*: Issue 25(6B): Freshwater Inflow: Science, Policy, and Management. organized by Montagna, P.A., M. Alber and P.H. Doering.

- Everglades National Park. 1997a. Florida Bay News, Spring 1997: Sea Turtles in Florida Bay. Retrieved 1/5/2004 from <http://www.nps.gov/ever/current/fbn97a-6.htm>
- Everglades National Park. 1997b. Mammals of Everglades National Park. List compiled by Robertson, W.B. and J. Kushlan. Retrieved 1/5/2004 from <http://www.nps.gov/ever/eco/mammals.htm>
- Everglades National Park. 1997c. Roseate Spoonbill. ENP, Homestead, FL. Available from: <http://www.nps.gov/ever/eco/spoonbil.htm>, 1p.
- Everglades National Park. 2004. Birds of Everglades National Park. Retrieved 1/5/2004 from <http://www.nps.gov/ever/eco/birds.htm>
- Fewster, R.M., S.T. Buckland, G.M. Siriwardena, S.R. Baillie and J.D. Wilson. 2000. Analysis of population trends for farmland birds using generalized additive models. *Ecology*, 81:1970–84.
- Finucane, J.H. and A. Dragovich, 1959. Counts of red tide organisms, *Gymnodinium breve*, and associated oceanographic data from Florida's west coast, 1954–1957. U.S. Fish Wildlife Service. *Special. Science Report, Fisheries*, 289:202–295.
- Fish, J.E. and M. Stewart. 1991. Hydrogeology of the surficial aquifer system of Dade County, Florida. U.S. Geological Survey Water Resources Investigation Report 90(4108):50.
- Florida Bay Program Management Committee. 2004. The Strategic Science Plan for Florida Bay. Management Committee for the Florida Bay and Adjacent Marine Systems Science Program. National Oceanic and Atmospheric Administration, Atlantic Oceanographic and Meteorological Laboratory, Miami, FL. 53 pp. http://www.aoml.noaa.gov/ocd/sferpm/PMC_SSP_FB_Final_Nov07_2004_HQ.pdf
- Florida Bay Science Program. 2003. A Synthesis of Research on Florida Bay. Florida Marine Research Institute (in preparation). Available from: http://www.aoml.noaa.gov/flbay/draft/wkn_contents.pdf
- Florida Fish and Wildlife Commission (FWC). 2004. Florida's Endangered Species, Threatened Species and Species of Special Concern. Florida Fish and Wildlife Commission, Tallahassee, FL (retrieved 1/29/2004) <http://myfwc.com/imperiled/species/pdf/Endangered-Threatened-Special-Concern-2004.pdf>
- Florida Museum of Natural History. 2004. Mangrove Life: Amphibians and Reptiles (retrieved 1/4/2004) <http://www.flmnh.ufl.edu/fish/southflorida/mangrove/lifereps.html>
- Fonseca, M.S. and J.S. Fisher. 1986. A comparison of canopy friction and sediment movement between four species of seagrasses with reference to their ecology and restoration. *Marine Ecology Progress Series*, 29:15–22.

- Fourqurean, J.W., J.C. Zieman and G.V.N. Powell. 1992. Phosphorus limitation of primary production in Florida Bay: evidence from the C:N:P ratios of the dominant seagrass *Thalassia testudinum*. *Limnology and Oceanography*, 37:162–71.
- Fourqurean, J.W. and M.B. Robblee. 1999. Florida Bay: a history of recent ecological changes. *Estuaries*, 22: 345–57.
- Fourqurean, J.W., J.N. Boyer, M.J. Durako, L.N. Hefty and B.J. Peterson. 2003a. Forecasting the response of seagrass distribution to changing water quality: statistical models from monitoring data. *Ecological Applications*, 13(2):474–89.
- Fourqurean, J.W., J.N. Boyer, M.J. Durako, L.N. Hefty and B.J. Peterson. 2003b. The Response of Seagrass Distribution to Changing Water Quality: Predictive Models from Monitoring Data. Joint Conference on the Science and Restoration of the Greater Everglades and Florida Bay Ecosystems. Palm Harbor, FL April 13–18, 2003.
- Fourqurean, J.W., M.J. Durako, M.O. Hall and L. N. Hefty. 2002. Seagrass Distribution in South Florida: A Multi-Agency-Coordinated Monitoring Program. In *The Everglades, Florida Bay, and Coral Reefs of the Florida Keys: An Ecosystem Sourcebook*, ed. J.W. Porter and K.W. Porter. CRC Press, Boca Raton, FL, 497–522 pp.
- Frankovich, T.A. and J.C. Zieman. 1994. Total epiphyte and epiphytic carbonate production on *Thalassia testudinum* across Florida Bay. *Bulletin of Marine Science* 54(3): 679–95.
- Frederick, P.C. and M.G. Spalding. 1994. Factors Affecting Reproductive Success of Wading Birds (Ciconiiformes) in the Everglades Ecosystem. In *Everglades: The Ecosystem and Its Restoration*, ed. S.M. Davis and J.C. Ogden. St. Lucie Press, Delray Beach, FL, 659–91 pp.
- Fry, B. and E.B. Sherr. 1984. $\delta^{13}\text{C}$ measurements as indicators of carbon flow in marine and freshwater ecosystems. *Contributions in Marine Science*, 27:13–47.
- Garcia, A.M. and J. P. Vierira. 1997. Anbundância e diversidade da assembléia de peixes dentro e fora de uma pradaria de *Ruppia maritima* L., no estuário da Lagoa dos Patos (RS, Brasil). *Atlantica* 19: 161-181.
- Gawlik, D.E. 2002. The effects of prey availability on the numerical response by wading birds. *Ecological Monographs*, 72:329–46.
- German, E.R. 2000. Regional Evaluation of Evapotranspiration in the Everglades. Water Resources Investigations Report 00-4217. U.S. Geological Survey, Tallahassee, FL.
- Gilmore, R.G. 1987. Fish, Macrocrustacean, and Avian Population Dynamics and Cohabitation in Tidally Influenced Impounded Subtropical Wetlands. In *Proceedings of a Symposium in Waterfowl and Wetland Management in the Coastal Zone of the*

- Atlantic Flyway, ed. W.R. Whitman and W.H. Meredith. Delaware Department of Natural Resources and Environmental Control, Dover, DE.
- Gleason, P.J. 1972. The Origin, Sedimentation, and Stratigraphy of a Calcite Mud Located in Southern Freshwater Everglades. Ph.D Dissertation, Pennsylvania State University, University Park, PA.
- Hall, M.O., J.J. Durako, J.W. Fourqurean and J.C. Zieman. 1999. Decadal changes in seagrass distribution and abundance in Florida Bay. *Estuaries* 22:445–59.
- Hanlon, R. and G. Voss. 1975. Guide to the Seagrasses of Florida, the Gulf of Mexico and the Caribbean Region. Sea Grant Field Guide 4, University of Miami Sea Grant, University of Miami, Miami, FL, 30 ff.
- Harrigan, P., J.C. Zieman and S.A. Macko. 1989. The base of nutritional support for the gray snapper (*Lutjanus griseus*): an evaluation based on a combined stomach content and stable isotope analysis. *Bulletin of Marine Science*, 44(1):565–77.
- Harrison, P.G. 1982. Seasonal and year-to-year variations in mixed intertidal populations of *Zostera japonica* and *Ruppia maritima*. *Aquatic Botany*, 14:357–71.
- Hartman, D.S. 1974. Distribution, Status, and Conservation of the Manatee in the United States. Technical Report NTIS PB81-140725, prepared for the U.S. Fish and Wildlife Service, National Fish and Wildlife Lab, 247 pp.
- Hastie, T. and R. Tibshirani. 1990. Generalized Additive Models. Chapman and Hall, Boca Raton, FL, 356 ff.
- Heald, E.J. 1969. The Production of Organic Detritus in a South Florida Estuary. Ph.D. dissertation, University of Miami, Coral Gables, FL. 110 ff.
- Heck, K.L., G. Hays and R.J. Orth. 2003. Critical evaluation of the nursery role hypothesis for seagrass meadows. *Marine Ecology Progress Series*, 253:123–36.
- Hicks, D.B. and L.A. Burns. 1975. Mangrove Metabolic Response to Alterations of Natural Freshwater Drainage to Southwestern Florida Estuaries. In Proceedings of International Symposium on Biology and Management of Mangroves, ed. Gerald Walsh *et al.* University of Florida, Gainesville, FL, 238–55 pp.
- Hitchcock, G., E. Philips and L. Brand. 2003. Chapter 6: Plankton Blooms. In A Synthesis of Research on Florida Bay, ed. W. Nuttle. Florida Bay Science Program, compiled for the Florida Bay Science Panel, U.S. Geological Survey, Florida Caribbean Science Center, Gainesville, FL, 276 p.
- Hittle *et al.* 2000

- Hittle, C., E. Patino and M. Zucker. 2001. Freshwater Flow from Estuarine Creeks into Northeastern Florida Bay: Water Resources Investigations Report 01-4164, US Geological Survey, Tallahassee, FL.
- Hoffmeister, J.E. 1974. Chapter 1: South Florida's Geologic Formations. In *Land from the Sea: The Geologic Story of South Florida*. University of Miami Press, Coral Gables, FL, 19–26 pp.
- Holmquist, J.F., G.V.N. Powell and S.M. Sogard. 1989a. Decapod and stomatopod communities of seagrass-covered mud banks in Florida Bay: inter- and intra-bank heterogeneity with special reference to isolated subenvironments. *Bulletin of Marine Science*, 44(1):251–62.
- Holmquist, J.F., G.V.N. Powell and S.M. Sogard. 1989b. Sediment, water level and water temperature characteristics of Florida Bay's grass-covered mud banks. *Bulletin of Marine Science*, 44(1):348–64
- Hopkins, A. 1986. The Development of the Overseas Highway. *Tequesta*, 46:45–58 pp.
- Iverson, R.L. and H.F. Bittaker. 1986. Seagrass distribution and abundance in eastern Gulf of Mexico coastal waters. *Estuarine, Coastal and Shelf Science*, 22:577–602.
- Jaap, W.C. 1984. The Ecology of South Florida Coral Reefs: A Community Profile: FWS/OBS 82/08. U.S. Field and Wildlife Service, Washington, D.C., 138 pp.
- Jewett-Smith, J. 1991. Factors influencing the standing crop of diatom epiphytes of the seagrass *Halodule wrightii* in south Texas seagrass beds. *Contributions in Marine Science*, 32:27–40.
- Johnson, D., J. Browder, D. Harper and S. Wong. 2002a. A Meta-Analysis and Synthesis of Existing Information on Higher Trophic Levels in Florida Bay: Final Report on Year One of a Two-Year Project IA5280-9-9031. Everglades National Park, Homestead, FL, and National Marine Fisheries Service, Miami, FL, 340 ff.
- Johnson, D., J. Browder, D. Harper and S. Wong. 2002b. A Meta-Analysis and Synthesis of Existing Information on Higher Trophic Levels in Florida Bay (model validation and prediction). Final Report on Year Two of a Two-Year Project IA5280-9-9031. Everglades National Park, Homestead, FL, and National Marine Fisheries Service, Miami, FL, 141 ff and appendices.
- Johnson, D.R., J.A. Browder and M.B. Robblee. 2004. Literature Review of Selected Organisms and Salinity-Related Responses from Florida Bay and Surrounding Areas. Final Report Prepared for the South Florida Water Management District, West Palm Beach, FL, 70 pp.
- Johnson, D.R., J.A. Browder and M.B. Robblee. 2005. Statistical Models of Florida Bay Fishes and Crustaceans to Evaluate Minimum Flow Levels in Florida Bay.

- Contribution No. PRD 04/05-06. Protected Resources and Biodiversity Division of the Southeast Fisheries Science Center, Miami, FL, 474 ff.
- Johnson, R.A. and R.J. Fennema. 1989. Conflicts over Flood Control and Wetland Preservation in the Taylor Slough and Eastern Panhandle Basins of Everglades National Park. In Symposium on Wetlands: Concerns and Success, ed. D.W. Fisk. American Water Resources Association, Bethesda, MD, 451–62 pp.
- Johnson, R.A. and T.J. VanLent. 1995 (June). Restoring Flows to the Shark Slough Basin, Everglades National Park. In Effects of Human-Induced Changes on Hydrologic Systems. American Water Resources Association, Bethesda, MD, 465–74 pp.
- Johnson, R.A., J.E. Wagner, D.J. Grigsby and V.A. Stern. 1988. Hydrologic Effects of the 1984 through 1986 L-31 Canal Drawdowns on the Northern Taylor Slough Basin of Everglades National Park: Technical Report SFRC-88/01. South Florida Research Center, Everglades National Park, Homestead, FL.
- Jurado, J.L. 2003. The Roles of Freshwater Discharge, Advective Processes and Silicon Cycling in the Development of Diatom Blooms in Coastal Waters of the Southwestern Florida Shelf and Northwestern Florida Bay. Ph.D. dissertation. Rosenstiel School of Marine and Atmospheric Sciences/University of Miami, Miami, FL.
- Kale, H.W., B. Pranty, B.M. Stith and C. W. Biggs. 1992. The Atlas of the Breeding Birds of Florida: Final Report. Florida Game and Fresh Water Fish Commission, Tallahassee, FL.
- Kale, H.W. and D.S. Maehr. 1990. Florida's Birds: A Handbook and Reference. Pineapple Press, Sarasota, FL.
- Kantrud, H.A. 1991. Wigeongrass (*Ruppia maritima*): A Literature Review. U.S. Fish and Wildlife Service, Fish and Wildlife Research 10. Northern Prairie Wildlife Research Center, Jamestown, ND. Retrieved 7/16/97 from: <http://www.npwrc.usgs.gov/resource/literatr/ruippia/ruippia.htm>.
- Kaplan, E.H. 1988. A Field Guide to Southeastern and Caribbean Seashores. New York: Houghton Mifflin Company.
- Kemp, W.M., R.R. Twilley, J.C. Stevenson, W. Boynton and J.C. Means. 1983. The decline of submerged vascular plants in the upper Chesapeake Bay: summary of results concerning possible causes. *Marine Technology Society Journal*, 1(17):78–89.
- Kenworthy, J. and A.C. Schwarzschild. 1995. Estimation of the Minimum Light Requirements of *Thalassia testudinum*, *Halodule wrightii*, and *Syringodium filiforme* in Biscayne Bay, FL. Report to South Florida Water Management District, West Palm Beach, FL.

- Koch, E.W. and C.J. Dawes. 1991. Influence of salinity and temperature on the germination of *Ruppia maritima* from the North Atlantic and Gulf of Mexico. *Aquatic Botany*, 40:387–91.
- Koch, M.S. 2003. High Salinity and Multiple Stressor Effects on Seagrass Communities of NE Florida Bay. Report (contract 12430) for South Florida Water Management District, West Palm Beach, FL, 30 ff.
- Koch, M.S. and J. M. Erskine. 2001. Sulfide as a phytotoxin to the tropical seagrass *Thalassia testudinum*: interactions with high salinity and temperature. *Journal of Experimental Marine Biology and Ecology*, 266:81–95.
- Koch, M.S. and M.J. Durako. 2004. High Salinity and Multiple Stressor Effects on Seagrass Communities of Northeast Florida Bay. Second Interim Report (contract 12430) for South Florida Water Management District, West Palm Beach, FL.
- Koch, M.S. and S.C. Snedaker. 1997. Factors influencing *Rhizophora mangle* (red mangrove) seedling development in Everglades carbonate soils. *Aquatic Botany*, 59:87–98.
- Koch, M.S., R.E. Benz and D.T. Rudnick. 2001. Solid-phase phosphorus pools in highly organic carbonate sediments of northeastern Florida Bay. *Estuarine, Coastal and Shelf Science*, 52:279–91.
- Kushan, J.A. 1988. Profiles: conservation and management of the American crocodile. *Environmental Management*, 12(6):777–90.
- Kushlan, J.D., O.L. Bass and L.C. McEwan. 1982. Wintering Waterfowl in Everglades National Park: Report T-670. Everglades National Park, South Florida Research Center, Homestead, FL.
- Kushlan, J.A. and F.J. Mazzotti. 1989. Historic and present distribution of the American crocodile in Florida. *Journal of Herpetology*, 23:1–7.
- Lavrentyev, P.J., H.A. Bootsma, T.H. Johengen, J.F. Cavaletto and W.S. Gardner. 1998. Microbial plankton response to resource limitation: insights from the community structure and seston stoichiometry in Florida Bay, USA. *Marine Ecology Progress Series*, 165:45–57.
- Lazar, A.C. and C.J. Dawes. 1991. A seasonal study of the seagrass *Ruppia maritima* in Tampa Bay, Florida: organic constituents and tolerances to salinity and temperature. *Botanica Marina*, 34:265–69.
- Lee, T.N., E. Williams, E. Johns, D. Wilson and N.P. Smith. 2002. Transport Processes Linking South Florida Coastal Ecosystems. In *The Everglades, Florida Bay, and Coral Reefs of the Florida Keys*, ed. J.W. Porter and K.G. Porter. CRC Press, Boca Raton, FL, 309–42 pp.

- Ley, J.A. 1992. Influence of Freshwater Flow on the Use of Mangrove Prop Root Habitat by Fish. PhD Dissertation, University of Florida, Gainesville, FL. 171 pp.
- Ley, J.A. and C.C. McIvor. 2002. Linkages between Estuarine and Reef Fish Assemblages: Enhancement by the Presence of Well-Developed Mangrove Shorelines. In *The Everglades, Florida Bay, and Coral Reefs of the Florida Keys: An Ecosystem Sourcebook*, ed. J.W. Porter and K.W. Porter. CRC Press, Boca Raton, FL 539–62 pp.
- Ley, J.A., C.C. McIvor and C.L. Montague. 1999. Fishes in mangrove prop-root habitats of northeastern Florida Bay: distinct assemblages across an estuarine gradient. *Estuarine, Coastal and Shelf Science*, 48:701–23.
- Light, S.S. and J.W. Dineen. 1994. Water Control in the Everglades: A Historical Perspective. In *Everglades: the Ecosystem and Its Restoration*, ed. S.M. Davis and J.C. Ogden. St. Lucie Press, Delray Beach, FL, 47–84 pp.
- Livingston, R.J. 1987. Historic Trends of Human Impacts on Seagrass Meadows of Florida. In *Florida Marine Research Publication 92: Proceedings of the Symposium on Subtropical-Tropical Seagrasses of the Southeastern United States*, ed. M.J. Durako, R.C. Phillips and R.R. Lewis. Florida Department of Natural Resources, Tallahassee, FL, 139–56 pp.
- Livingston, R.J. 1989. Ecosystem research and resource management: application to Florida Bay. *Bulletin of Marine Science*, 44(1):517–18 (abstract).
- Lorenz, J.J. 1999. The response of fishes to physicochemical changes in the mangroves of northeast Florida Bay. *Estuaries*, 22:500–17.
- Lorenz, J.J. 2000. Impacts of Water Management on Roseate Spoonbills and Their Piscine Prey in the Coastal Wetlands of Florida Bay. Doctoral Dissertation submitted to University of Miami, Coral Gables, FL.
- Lorenz, J.J., J.C. Ogden, R.D. Bjork and G.V.N. Powell. 2002. Nesting Patterns of Roseate Spoonbills in Florida Bay 1935–1999: Implications of Landscape-Scale Anthropogenic Impacts. In *The Everglades, Florida Bay, and Coral Reefs of the Florida Keys: An Ecosystem Sourcebook*, ed. J.W. Porter and K.W. Porter. CRC Press, Boca Raton, FL, 563–606 pp.
- Madden, C.J. and A.M. McDonald. 2004 (December). Analysis of Salinity Conditions Impacting the SAV Community. In support of minimum flows and levels for Florida Bay. Internal Memorandum, Coastal Ecosystems Department, South Florida Water Management District, West palm Beach, FL.
- Madden, C.J., A.M. McDonald, M.J. Hunt, W.M. Kemp and D. Gruber. 2003 (May). Summary: Ecosystem Process Models of Seagrass Communities in Florida Bay: Internal Memorandum. South Florida Water Management District, West Palm Beach, FL.

- Marshall, F.E., D. Smith and D. Nickerson. 2004. Using Statistical Models to Simulate Salinity Variation and Other Physical Parameters in North Florida Bay. Cetacean Logic Foundation, Inc. New Smyrna Beach, FL, 36 pp.
- Marshall, F.E. 2005 (July 11). Extension of Taylor River Salinity Historical Reconstruction. A letter report prepared for the South Florida Water Management District (contract C-15975-WO05), West Palm Beach, FL, by Environmental Consulting and Technology, Inc., New Smyrna Beach, FL
- Marshall, F.E. 2005 (June 27). Monthly Salinity Simulations for Taylor River Interim CERP Update Runs. A letter report prepared for the South Florida Water Management District (contract C-15975-WO05), West Palm Beach, FL, by Environmental Consulting and Technology, Inc., New Smyrna Beach, FL.
- Marshall, F.E. 2004a. Taylor River Salinity Historical Reconstruction. A letter report prepared for the South Florida Water Management District (contract C-15975-WO05), West Palm Beach, FL, by Environmental Consulting and Technology, Inc., New Smyrna Beach, FL
- Marshall, F.E. 2004b. ICU Runs (draft). Summary Report for the Southern Estuaries Sub-Team of [CERP] Regional Evaluation Team. Environmental Consulting and Technology, Inc. New Smyrna Beach, FL.
- Matheson Jr., R.E., D.K. Camp, S.M. Sogard and K.A. Bjorgo. 1999. Changes in seagrass-associated fish and crustacean communities on Florida Bay mud banks: the effects of recent ecosystem changes? *Estuaries*, 22:534–51.
- Mazzotti, F.J. and W.A. Dunson. 1984. Adaptations of *Crocodylus acutus* and alligator for life in saline water. *Comparative Biochemistry. Physiology*, 79A(4):641–46.
- Mazzotti, F.J. 1999. The American crocodile in Florida Bay. *Estuaries*, 22:552–61.
- Mazzotti, F.J. 1989. Factors affecting the nesting success of the American crocodile, *Crocodylus acutus*, in Florida Bay. *Bulletin of Marine Science*, 44:220–28.
- Mazzotti, F.J., B. Bohnsack, M.P. McMahon and J.R. Wilcox. 1986. Field and laboratory observations on the effects of high temperature and salinity on hatchling *Crocodylus acutus*. *Herpetologica*, 42(2):191–96.
- Mazzotti, F.J., J.A. Kushlan, A. Dunbar-Cooper. 1988. Desiccation and cryptic nest flooding as probable causes of egg mortality in the American crocodile, *Crocodylus acutus*, in Everglades National Park, Florida. *Florida Scientist*, 51(2):65–72.
- McIvor, C.C., J.A. Ley and R.D. Bjork. 1994. Changes in Freshwater Inflow from the Everglades to Florida Bay, Including Effects on Biota and Biotic Processes: A Review. In *Everglades: The Ecosystem and Its Restoration*, ed. S.M. Davis and J.C. Ogden. St. Lucie Press, Delray Beach, FL, 117–46 pp.

- McMillan, C. 1974. Salt Tolerance of Mangroves and Submerged Aquatic Plants. In Ecology of Halophytes, ed. R.J. Reimold and W.H. Queen. Academic Press, New York, NY, 379–90 pp.
- McMillan, C. and F.N. Moseley. 1967. Salinity tolerances of five marine spermatophytes of Redfish Bay, Texas. *Ecology*, 48:503–06.
- Meeder, J.F. and L.B. Meeder. 1989. Hurricanes in Florida Bay: a dominant physical process. *Bulletin of Marine Science*, 44(1):518 (abstract).
- Meyer, F.W. 1989. Hydrogeology, Groundwater Movement, and Subsurface Storage in the Floridan Aquifer System in Southern Florida. Professional Paper 1403-G, U.S. Geological Survey, 59 ff.
- Micheli, F. and C.H. Peterson. 1999. Estuarine vegetated habitats as corridors for predator movements. *Conservation Biology*, 13:869–81.
- Moler, P.E. 1991. American Crocodile Population Dynamics. Final Report to Study 7532, Florida Game and Fresh Water Fish Commission, Bureau of Wildlife Research, Tallahassee, FL.
- Montague, C.L. and J.A. Ley. 1993. A possible effect of salinity fluctuation on abundance of benthic vegetation and associated fauna in northeastern Florida Bay. *Estuaries*, 16:703–17.
- Montague, C.L., R.D. Bartleson and J.A. Ley. 1989. Assessment of Benthic Communities along Salinity Gradients in Northeastern Florida Bay. Final Report to the South Florida Research Center, Everglades National Park, Homestead, FL.
- Montague, C.L., E. Chipouras and L.M. Jones. 1998. Responses of Submerged Macrophytes to Freshwater Inflow to Florida Bay. Prepared for the South Florida Water Management District, West Palm Beach, FL.
- Morrison, D. and D.L. Bean. 1997. Benthic Macrophyte and Invertebrate Distribution and Seasonality in the Everglades–Florida Bay Ecotone. Final Project Report to South Florida Research Center, Everglades National Park, Homestead, FL.
- Murphy, L.R., S.T. Kinsey and M.J. Durako. 2003. Physiological effects of short-term salinity changes on *Ruppia maritima*. *Aquatic Botany*, 75:293–309.
- Musick, J.A., M.M. Harbin, S.A. Berkeley, G.H. Burgess, A.M. Eklund, L. Findley, R.G. Gilmore, J.T. Golden, D.S. Ha, G.R. Huntsman, J.C. McGovern, S.J. Parker, S.G. Poss, E. Sala, T.W. Schmidt, G.R. Sedberry, H. Weeks and S.G. Wright. 2000. Marine, estuarine, and diadromous fish stocks at risk of extinction in North America. *Fisheries*, 25(11):6–30.
- Myers, R.L. 1986. Florida's freezes: an analog of short-duration nuclear winter events in the tropics. *Florida Scientist*, 49:104–15.

- Myers, R.L. and J. Ewel. 1990. Ecosystems of Florida. University of Central Florida Press. Orlando, FL.
- National Marine Fisheries Service. 2000. Status Review of Smalltooth Sawfish (*Pristis pectinata*). National Oceanic and Atmospheric Administration (NOAA), NMFS, Silver Spring, MD, 73 ff. http://www.nmfs.noaa.gov/pr/readingrm/statrvws/Smalltooth_sawfish.PDF
- National Oceanic and Atmospheric Administration 2000. Florida Bay Zones. Florida Bay and Adjacent Marine Systems. Atlantic Oceanographic and Meteorological Laboratory, Miami, FL. Available from Science Programs website: <http://www.aoml.noaa.gov/ocd/sferpm/sub.html>
- Nelsen JE, Ginsburg RN. Calcium carbonate production by epibionts on *Thalassia* in Florida Bay. *J Sediment Petrol* 1986;56:622—8.
- Nuttle, W.K., J.W. Fourqwean, B.J. Cosby, J.C. Zieman and M.B. Robblee. 2000. Influence of net freshwater supply on salinity in Florida Bay. *Water Resources Research*, 36:1805–22.
- Nuttle, W.K., R. Price and P.K. Swart. 2001. Estimating Evaporation from Florida Bay. Estuarine Research Federation, Sixteenth Biennial Conference, November 2001, St. Petersburg Beach, FL. (abstract, p. 102).
- Nuttle, W.K., R. Price and P.K. Swart. 2003. Variation of Evaporation from Florida Bay. Joint Conference on the Science and Restoration of the Greater Everglades and Florida Bay Ecosystem, April 2003, Palm Harbor, FL (program and abstracts, pp 61–63).
- Nuttle, W.K. and R.S. Teed. 2002. Wetland Hydrology and Estuarine Salinity Models for the Taylor Slough/C-111 Area: Version 1. Report prepared for the Everglades National Park by the Cadmus Group, Inc., Watertown, MA.
- Oberhofer, L. and S. Bass. 2003. Everglades National Park. In South Florida Wading Bird Report (volume 9), ed. G.E. Crozier and D.E. Gawlik. South Florida Water Management District, West Palm Beach, FL, 8 ff.
- Odum, W.E. 1970. Pathways of Energy Flow in a South Florida Estuary. Ph.D. dissertation, University of Miami, Coral Gables, FL, 162 ff.
- Odum, W.E., C.C. McIvor and T.J. Smith. 1982. The Ecology of the Mangroves of South Florida: A Community Profile. FWS/OBS-82/24. U.S. Fish and Wildlife Service, Office of Biological Services, Washington, D.C., 144 pp.
- Ogden, J.C. 1978a. Status and nesting biology of the American crocodile, *Crocodylus acutus*, in Florida. *Journal of Herpetology*, 12(2):183–96.

- Ogden, J.C. 1978b. American Crocodile. In Amphibians and Reptiles (volume 3 of Rare and Endangered Biota of Florida), ed. R.W. McDiarmid. University Presses of Florida, Gainesville, FL, 21–22 pp.
- Olmsted, I.G., L.L. Lloyd and R.P. Russell. 1981. Vegetation of the Southern Coastal Region of Everglades National Park between Flamingo and Joe Bay. Report T-620. National Park Service, South Florida Research Center, Homestead, FL 18 pp.
- Orlando, S.P., M.B. Robblee and C.J. Klein. 1997. Salinity Characteristics of Florida Bay: A Review of the Archived Data Set (1955–1995). National Oceanic and Atmospheric Administration, Office of Ocean Resources Conservation and Assessment, Silver Spring, MD, 38 pp.
- Parker, G.G., G.E. Ferguson, S.K. Love *et al.* 1955. Water Resources of Southeastern Florida. USGS Paper 1255. U.S. Geological Survey, Water Supply, 965 pp.
- Paul, R. 2003 (January). Roseate Spoonbill. In Florida's Breeding Bird Atlas: A Collaborative Study of Florida's Birdlife. Florida Fish and Wildlife Conservation Commission. Retrieved 9/19/2005 from: <http://www.myfwc.com/bba/>
- Perkins, R.D. 1977. Depositional Framework of Pleistocene Rocks in South Florida. In: P. Enos and R.D. Perkins (Editors), Quaternary Sedimentation in South Florida. Geological Society of America, Memoir 147, pp131–198.
- Perkins, W.A. and T.K. MacVicar. 1991. A Computer Model to Simulate Natural South Florida Hydrology. Draft report. South Florida Water Management District, West Palm Beach, FL.
- Phillips, R.C. 1960. Observations on the Ecology and Distribution of the Florida Seagrasses. Professional Papers Series 2. Florida Board of Conservation, 1–61 pp.
- Phlips, E.J. and S. Badylac. 1996. Spatial variability in phytoplankton standing stock and composition in a shallow inner-shelf lagoon of Florida Bay, Florida. *Bulletin of Marine Science*, 58:203–16.
- Phlips, E.J., S. Badylak and T. Grosskopf. 2002. Factors affecting the abundance of phytoplankton in a restricted subtropical lagoon: the Indian River Lagoon, Florida, USA. *Estuarine, Coastal and Shelf Science*, 55:385–402.
- Phlips, E., C. Zeman and P. Hansen. 1989. Growth, photosynthesis, nitrogen fixation and carbohydrate production by a unicellular cyanobacterium, *Synechococcus* sp. (Cyanophyta). *Journal of Applied Physiology*, 1:137–45.
- Phlips, E.J., S. Badylak and T.C. Lynch. 1999. Blooms of the picoplanktonic cyanobacterium *Synechococcus* in Florida Bay, a subtropical inner-shelf lagoon. *Limnology and Oceanography*, 44:1166–75.

- Poulakis, G. R. and J. C. Seitz. 2004. Recent occurrence of the smalltooth sawfish, *Pristis pectinata* (Elasmobranchiomorphi: Pristidae), in Florida Bay and the Florida Keys, with comments on sawfish ecology. *Florida Scientist*, 67(1):27–35.
- Powell, A.B., D.E. Hoss, W.F. Hettler, D.S. Peters and S. Wagner. 1989. Abundance and distribution of ichthyoplankton in Florida Bay and adjacent waters. *Bulletin of Marine Science*, 44:35–48.
- Powell, A.B. 2003. Larval abundance, distribution, and spawning habits of spotted seatrout (*Cynoscion nebulosus*) in Florida Bay, Everglades National Park, Florida. *Fishery Bulletin*, 101:704–11.
- Powell, J.A. and G.B. Rathbun. 1984. Distribution and abundance of manatees along the northern coast of the Gulf of Mexico. *Northeast Gulf Science*, 7(1):1–28.
- Prager, Ellen, and Bob Halley. 1997. Sediment Bottom Map for Florida Bay. Open File Report #97-526. US Geological Survey.
- Price, R. 2001. Geochemical Determinations of Groundwater Flow in Everglades National Park. Ph.D. dissertation. University of Miami, Coral Gables, FL, 307 ff.
- Redfield, A.C. 1958. The biological control of chemical factors in the environment. *American Scientist*, 46: 205–22.
- Redfield, A. C. 1967. Postglacial change in sea level in the western North Atlantic Ocean. *Science*, 157:687–92
- Reese, R.S. and K.J. Cunningham. 2000, Hydrogeology of the Gray Limestone Aquifer in Southern Florida: U.S. Geological Survey Water-Resources Investigations Report 99-4213.
- Robblee, M.B., S.D. Jewell and T.S. Schmidt. 1991. Temporal and Spatial Variation in the Pink Shrimp, *Penaeus duorarum*, in Florida Bay and Adjacent Waters of Everglades National Park. In SFRC 1991 Annual Reports, ed. R.F. Doren and C. Dofferrmyre. South Florida Research Center, Everglades National Park, Homestead, FL.
- Robblee, M.B., T.R. Barber, P.R. Carlson Jr., M.J. Durako, J.W. Fourqurean, L.K. Muehlstein, D. Porter, L.A. Yarbro, J.C. Zieman and R.T. Zieman. 1991. Mass mortality of the tropical seagrass *Thalassia testudinum* in Florida Bay (USA). *Marine Ecology Progress Series*, 71:297–99.
- Ross, C.A. and W.E. Magnusson. 1989. Living Crocodilians. In Crocodiles and Alligators, ed. C.A. Ross. Facts on File, Inc., New York, NY, 58–75 pp.
- Ross, M.S., E.E. Gaiser, J.F. Meeder and M.T. Lewin. 2002. Multi-Taxon Analysis of the “White Zone”: A Common Ecotonal Feature of the South Florida Coastal Wetlands. In The Everglades, Florida Bay, and Coral Reefs of the Florida Keys: An Ecosystem

- Sourcebook, ed. J.W. Porter and K.W. Porter. CRC Press, Boca Raton, FL, 205–238 pp.
- Ross, M.S., J.F. Meeder, J.P. Sah, P.L. Ruiz and G.J. Telesnicki. 2000. The southeast saline Everglades revisited: 50 years of coastal vegetation change. *Journal of Vegetation Science*, 11:101–12.
- Rowe, C.L. and W.A. Dunson. 1995. Individual and interactive effects of salinity and initial fish density on a salt marsh assemblage. *Marine Ecology Progress Series*, 128:271–78.
- Rudnick, D. 2004 (January). Appendix A.5: Florida Bay Conceptual Ecological Model. In Comprehensive Everglades Restoration Plan, Part 1: Monitoring and Supporting Research. United States Army Corps of Engineers, Jacksonville, FL, and South Florida Water Management District, West Palm Beach, FL, A99–115 pp.
- Rudnick, D.T., P.B. Ortner, J.A. Browder, and S.M. Davis. 2005. A conceptual model of Florida Bay. *Wetlands* 25:870–883.
- Rudnick, D.T., Z. Chen, D.L. Childers, J.N. Boyer and T.D. Fontaine III. 1999. Phosphorus and nitrogen inputs to Florida Bay: the importance of the Everglades watershed. *Estuaries*, 22: 398–416.
- Russell, R.P., L.L. Lloyd and T.C. Olmstead. 1981. Vegetation of the Southern Coastal Region of Everglades National Park between Flamingo and Joe Bay. Report T-620. National Park Service, South Florida Research Center. Homestead, FL.
- Rutherford, E.S., T.W. Schmidt, and J.T. Tilmant. 1986. Early Life History of Spotted Seatrout, Red Drum, Gray Snapper, and Snook in Everglades National Park, Florida. Rept 86/07. Homestead, FL. Everglades N.P., South Florida Research Center. 99pp.
- Rutherford, E.S., J.T. Tilmant, E.B. Thue and T.W. Schmidt. 1989. Fishery harvest and population dynamics of spotted seatrout (*Cynoscion nebulosus*) in Florida Bay and adjacent waters. *Bulletin of Marine Science*, 44(1):108–25.
- Schmidt, T.W. 1979. Ecological Study of Fishes and Water Quality Characteristics of Florida Bay. Report RSP-EVER N-36. Everglades National Park, National Park Service, South Florida Research Center, Homestead, FL, 145 ff.
- Schomer, N.S. and R.D. Drew. 1982. An Ecological Characterization of the Lower Everglades, Florida Bay, and the Florida Keys. WS/OBS-82/58.1. U.S. Fish and Wildlife Service, Office of Biological Services, Washington, D.C.
- Seeliger, U., C. Cordazzo and E.W. Koch. 1983. Germination and algal-free laboratory culture of wigeongrass, *Ruppia maritima*. *Estuaries*, 7:176–78.

- Shaw, R., W. Wiseman Jr., R.E. Turner, L.J. Rouse Jr., R.E. Condrey and F.J. Kelly Jr. 1985. Transport of larval gulf menhaden, *Brevoortia patronus*, in continental shelf waters of western Louisiana: a hypothesis. *American Fisheries Society*, 114:452–60.
- Sklar, F., C. McVoy, R. VanZee, D.E. Gawlik, K. Tarboton, D. Rudnick and S. Miao. 2002. The Effects of Altered Hydrology on the Ecology of the Everglades. In *The Everglades, Florida Bay, and Coral Reefs of the Florida Keys: An Ecosystem Sourcebook*, ed. J.W. Porter and K.W. Porter. CRC Press, Boca Raton, FL, 39–82 pp.
- Sklar, F.H. and J.A. Browder. 1998. Coastal environmental impacts brought about by alterations to freshwater flow in the Gulf of Mexico. *Environmental Management*, 22:547–62.
- Smith, D. 2001. Marine Physical Conditions: Everglades National Park October 1998 through December 1999. South Florida Natural Resources Technical Report 2001–2002. Everglades National Park, Homestead, FL
- Snedaker, S. 1989. Overview of ecology of mangroves and information needs for Florida Bay. *Bulletin of Marine Science*, 44: 341–47.
- Snow, R.W. 1991. The Distribution and Relative Abundance of the Florida Manatee in Everglades National Park. Annual Report. South Florida Research Center, Everglades National Park, Homestead, FL, 26 pp.
- Sogard, S.M., G.V. Powell and J.G. Holmquist. 1989a. Spatial distribution and trends in abundance of fishes residing in seagrass meadows on Florida Bay mud banks. *Bulletin of Marine Science*, 44:179–99.
- Sogard, S.M., G.V. Powell and J.G. Holmquist. 1989b. Utilization by fishes of shallow, seagrass-covered banks in Florida Bay: species composition and spatial heterogeneity. *Environmental Biology of Fishes*, 24:53–65.
- South Florida Water Management District (SFWMD). 1992a. Surface Water Improvement and Management Plan for the Everglades. West Palm Beach, FL, 471 pp.
- South Florida Water Management District. 1992b. *A Primer to the South Florida Water Management Model (Version 3.5)*. Hydrologic Systems Modeling Division, West Palm Beach, FL, 233 pp.
- South Florida Water Management District. 2000. Lower East Coast Regional Water Supply Plan. Prepared by Water Supply Planning and Development Department, West Palm Beach, FL.
- South Florida Water Management District. 2002. MFLs for the Northwest Fork of the Loxahatchee River. Final Draft.

- South Florida Water Management District. 2003. Selected Methodology for Long-Term (1965–2000) Solar Radiation and Potential Evapotranspiration Estimation for the SFWMM 2000 Update. Internal Memorandum to Ken Tarboton from Michelle Irizarry-Ortiz, December 12, 2003. South Florida Water Management District, West Palm Beach, FL.
- Steidinger, K., W. Richardson, M.B. Neely, G. McRae, S. Richards, R. Bray, T.H. Perkins and C.R. Tomas. 2001. Florida Bay Microalgal Blooms (abstract). Florida Bay Science Conference, Key Largo, FL.
- Stockman, K.W., R.N. Ginsburg and E.A. Shinn. 1967. The production of lime mud by algae in south Florida. *Journal of Sedimentary Petrology*, 37(2):633–48.
- Stoner, A.W. 1983. Distribution of fishes in seagrass meadows: role of macrophyte biomass and species composition. *Fisheries Bulletin*, 81(4):837–46.
- Stumpf, R.P., M.L. Frayer, M.J. Durako and J.C. Brock. 1999. Variations in water clarity and bottom albedo in Florida Bay from 1985 to 1997. *Estuaries*, 22:431–44.
- Sutula, M., J.W. Day, J. Cable and D. Rudnick. 2001. Hydrological and nutrient budgets of freshwater and estuarine wetlands of Taylor Slough in Southern Everglades, Florida (U.S.A). *Biogeochemistry*, 56:287–310.
- Swartzman, G., C. Huang and S. Kaluzny. 1992. Spatial analysis of Bering Sea groundfish survey data using generalized linear models. *Canadian Journal of Fisheries and Aquatic Sciences*, 49:1366–78.
- Tabb, D. C. and R. B. Manning. 1961. A checklist of the flora and fauna of northern Florida Bay and adjacent brackish waters of the Florida mainland collected during the period July 1957 through September 1960. *Bulletin Marine Science Gulf and Caribbean*, 11(4):552–649.
- Tabb, D.C. and D.L. Dubrow. 1962. Biological Data on Pink Shrimp, *Penaeus duorarum*, of North Florida Bay and Adjacent Estuaries in Monroe County, Florida, September 1957–March 1962. Report ML 62239. Institute of Marine Science, University of Miami, Coral Gables, FL.
- Tabb, D.C. and M.A. Roessler, 1989. History of studies of juvenile fishes of coastal waters of Everglades National Park. *Bulletin of Marine Science*, 44(1):23–34
- Tabb, D.C., D.L. Dubrow, and R.B. Manning. 1962. The ecology of north Florida Bay and adjacent estuaries. Florida Board of Conservation Technical Series 39:1–81.
- Thayer, G.W. and A.J. Chester. 1989. Distribution and abundance of fishes among basin and channel habitats in Florida Bay. *Bulletin of Marine Science*, 44:200–19.

- Thayer, G.W., A.B. Powell and D.E. Hoss. 1999. Composition of larval, juvenile, and small adult fishes relative to changes in environmental conditions in Florida Bay. *Estuaries*, 22:518–33.
- Thayer, G.W., W.J. Kenworthy and M.S. Fonseca. 1984. The Ecology of Eelgrass Meadows of the Atlantic Coast: A Community Profile. FWS/OBS-84/02. United States Fish and Wildlife Service, United States Department of the Interior, Washington, D.C.
- Tilmant, J.T. 1989. A history and overview of recent trends in the fisheries in Florida Bay. *Bulletin of Marine Science*, 44(1):3–22.
- Tomasko, D.A., C.J. Dawes and M.O. Hall. 1996. The effects of anthropogenic nutrient enrichment on turtle grass (*Thalassia testudinum*) in Sarasota Bay, Florida. *Estuaries*, 19(2B):448–56.
- Trappe, C.A. and G.A. Brewster-Wingard. 2001. Molluscan Fauna from Core 25B, Whipray Basin, Central Florida Bay, Everglades National Park. USGS Open File Report 01-143. U.S. Geological Survey.
- Trimble, P.J., E.R. Santee, R. Van Zee, L.G. Cadavid, J.T.B. Obeysekera and A. Ali. 2001. Simulation of the Influence of Climatic Variability on Freshwater Inflow to Florida Bay during the 20th Century (abstract). Florida Bay Science Conference, Key Largo, FL.
- Tucker, M.E. and V.P. Wright. 1990. Carbonate Sedimentology. Blackwell, Oxford, 482 pp.
- Turney, W.J. and B.F. Perkins. 1972. Sedimenta III: Molluscan Distribution in Florida Bay. Comparative Sedimentology Lab, University of Miami, Rosenstiel School of Marine and Atmospheric Science, Miami, FL. 37pp.
- Twilley, R., W. Kemp, K. Stayer, J.C. Stevenson and W. Boynton. 1985. Nutrient enrichment of estuarine submersed vascular plant communities. *Marine Ecology Progress Series*, 23:179–91.
- Tyler-Walters, H. 2001. *Ruppia maritima*. Beaked Tasselweed. Marine Life Information Network: Biology and Sensitivity Key Information Sub- Programme (online). Plymouth: Marine Biological Association of the United Kingdom [cited 02/14/2005]. <http://www.marlin.ac.uk/species/Ruppiamaritima.htm>
- U. S. Environmental Protection Agency. 1987. Estuary Program Primer. EPA Office of Marine and Estuarine Protection. Washington, D.C.
- U.S. Fish and Wildlife Service. 1998. Multi-Species Recovery Plan for the Threatened and Endangered Species of South Florida (volume 1 of 2): The Species. Technical/Agency Draft. Vero Beach, FL.

- United States Army Corps of Engineers (USACE). 1999. Central and Southern Florida Project Comprehensive Review Study, Final Integrated Feasibility Report and Programmatic Environmental Impact Statement. USACE, Jacksonville, FL.
- Van Lent, T.J., R. Johnson and R. Fennema. 1993. Water Management in Taylor Slough and Effects on Florida Bay. Technical Report SRFC 93-03. Everglades National Park, Homestead, FL.
- Verhoeven, J.T.A. 1975. *Ruppia* communities in the Camarque, France: distribution and structure in relation to salinity and salinity functions. *Aquatic Botany*, 1:217–41
- Verhoeven, J.T.A. 1979. The ecology of *Ruppia*-dominated communities in western Europe: distribution of *Ruppia* representatives in relation to their autecology. *Aquatic Botany*, 6:197–268.
- Verhoeven, J.T.A. 1980b. The ecology of *Ruppia*-dominated communities in western Europe: aspects of production, consumption and decomposition. *Aquatic Botany*, 8:209–53.
- Verhoeven, J.T.A. and W. van Vierssen. 1978. Distribution and structure of communities dominated by *Ruppia*, *Zostera* and *Potamogeton* species in the inland waters of “De Bol,” Texel, Netherlands. *Estuarine and Coastal Marine Science*, 6:417–28.
- Verhoeven, J.T.A. 1980a. The ecology of *Ruppia*-dominated communities in western Europe: synecological classification, structure and dynamics of the macroflora and macrofaunal communities. *Aquatic Botany*, 8:185.
- Virnstien, R.W. and K.D. Cairns. 1986. Seagrass Maps of the Indian River Lagoon. Final Report. Prepared for the St. Johns River Water Management District, Palatka, FL, by Seagrass Ecosystems Analysts, Inc., Vero Beach, FL.
- Virnstien, R.W., P.S. Mickelson, K.D. Cairns and M.A. Capone. 1983. Seagrass beds versus sand bottoms: the trophic importance of their associated benthic invertebrates. *Florida Scientist*, 46: 363–81.
- Wanless, H. R., R. W. Parkinson, and L. P. Tedesco. 1994. Sea level control on stability of Everglades wetlands. p. 199-223. In S. M. Davis and J.C. Ogden (eds.) Everglades: the Ecosystem and Its Restoration. St. Lucie Press, Delray Beach, FL, USA.
- Wanless, H.R. and M.G. Taggett. 1989. Origin, growth and evolution of carbonate mud banks in Florida Bay. *Bulletin of Marine Science*, 44(1):454–89.
- White, W.A. 1970. The geomorphology of the Florida peninsula. *Florida. Bureau of Geology Bulletin*, 51.
- Wingard, G.L., T.M. Cronin, G.S. Dwyer, S.E. Ishman, D.A. Willard, C.W. Holmes, C.E. Bernhardt, C.P. Williams, M.E. Marot, J.B. Murray, R.G. Stamm, J.H. Murray and C. Budet. 2003, Ecosystem History of Southern and Central Biscayne Bay: Summary

- Report on Sediment Core Analyses. USGS OFR 03-375. U.S. Geological Survey, 110 pp.
- Woodward-Clyde, Inc. 1998. St. Lucie Estuary Historical, SAV, and American Oyster Literature Review. Prepared for the South Florida Water Management District, West Palm Beach, FL.
- Zieman, J.C. 1982. A Community Profile: The Ecology of the Seagrass Ecosystem of South Florida. FWS/OBS-82/25. U.S. Fish and Wildlife Service, Office of Biological Services. Washington, D.C., 123 ff.
- Zieman, J.C., R. Orth, R.C. Phillips, G. Thayer and A. Thorhaug. 1984. The Effects of Oil on Seagrass Ecosystems. In *Restoration of Habitats Impacted by Oil Spills*, ed. J. Cairns and A.L. Buikema Jr. Butterworth, Boston, MA, 37–64 pp.
- Zieman, J.C., J.W. Fourqurean, and R.L. Iverson. 1989. Distribution, abundance and productivity of seagrasses and macroalgae in Florida Bay. *Bulletin of Marine Science* 44: 292–311.
- Zieman, J.C. and R.T. Zieman. 1989. The Ecology of the Seagrass Meadows of the West Coast of Florida: A Community Profile. Biological Report 85, U.S. Fish and Wildlife Service, United States Department of the Interior, Washington, D.C.
- Zieman, J.C., J.W. Fourqurean and T.A. Frankovich. 1999. Seagrass die-off in Florida Bay: long-term trends in abundance and growth of turtle grass, *Thalassia testudinum*. *Estuaries*, 22:460–70.